



Periodontal Therapy Home Care Instructions

A periodontal therapy program is a “conservative program of 3, 4, or 6 visits to control periodontal (gum) disease.” By “conservative” we mean non-surgical. Periodontal (gum) disease is not curable – but it is controllable.

Periodontal therapy aids in the removal of the bacterial infection of the hard and soft tissue which supports the teeth. Left untreated, the bacterial infection can cause the bone and surrounding tissues which support the teeth to deteriorate. Eventually teeth become loose and may fall out.

Common signs and symptoms of periodontal (gum) disease are:

- Bleeding gums
- Sore or swollen gums
- Pockets (bone loss with infection) in gum tissue greater than 3 mm
- Loosening teeth
- Malodor (bad breath)

The severity of the periodontal disease (mild, moderate or severe) determines the length of your program. Our goal is to eliminate bleeding points and to decrease pocket depth in order to prevent further loosening or loss of teeth.

As dental professionals, we will monitor your oral health and assist you in the control of plaque, calculus and bacteria. Remember to keep your recommended periodontal maintenance appointments, typically between every 6 weeks and 4 months. Bacteria that is associated with gum infection and attachment loss will reoccur within the pockets in just 6-12 weeks, although individual results may vary.

- If soreness or tenderness occurs, gently rinse your mouth 2-3 times per day with warm salt water (one tsp salt/8 oz. water). Start immediately when you get home.
- Your gums may become “achy”. You may also use Advil, Motrin, or Ibuprofen as directed.
- Your next meal should be soft. Avoid any hard or “chippy” foods like potato chips, popcorn, Fritos, etc. for the next 3-4 days.
- Swelling or jaw stiffness occur very rarely, however if it does, place warm moist towels to the face in the area of stiffness.
- Twice a day (once after breakfast and once before bedtime), brush gently along the gum line using a soft manual toothbrush or electric toothbrush for 2 minutes.
- Floss (remember to wrap floss around tooth and go below the gum line, use floss holders for hard to reach areas). Scrape up and down at least 3 times per area.
- Apply Peridex onto the brush or floss (when or if your hygienist recommends it).
- Rinse with Peridex mouthwash for 30 seconds (when or if your hygienist recommends it).

∞ PERSONALIZED AND COMFORTABLE ∞

KEITH N. WARR DDS, 1060 EAST 100 SO., #201, SALT LAKE CITY, 84102 Phone: 801-355-5385

